| INFORMATION DISCLOSURE STATEMENT BY APPLICANT (use as many sheets as necessar) 27 Sheet 5 of 9 Aktromey Docket No. 98,385-1 OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pages, Etc). OTHER DOCUMENTS (including Partie), Title, Date, Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). Attemption of Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). Attemption Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). It including Pages, Etc). OTHER DOCUMENTS (including Pages, Etc). It including Including Pages, Etc). It including Including Pages, Etc). It including Includi | Substitute fo | r form 1449A | PTO | | plete if Known | | | | | | | |
|---|---------------|--------------|---|--|--|--|--|--|--|--|--|--|
| STATEMENT BY APPLICANT (use as many sheets as necessant) (use as many sheets as necessary said of the speed of | | | | | Application No. | 09/898,234 | | | | | | |
| STATEMENT BY APPLICAN (use as many sheets as necessary (use as near was use many sheets as necessary (use as near was use many sheets as necessary (use as near was use many sheets as necessary (use as necessary (use as near was use many sheets as necessary (use as nec | | INFOR | MATION DISCI | OSURF | Filing Date: | 7-3-01 | | | | | | |
| (use as many sheets as necessary 27 | | | | | First Named Inventor | Hauptmann et al., | | | | | | |
| OTHER DOCUMENTS (Including Active, Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (Including Active, Title, Date, Pertinent Pages, Etc). Evans, "The steroid and thyroid hormone receptor superfamily," Science 240:889-95 (1988). Frohman et al., "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer," Proc. Natl. Acad. Sci. U. S. A. 85:8988-9002 (1988). Satanaga et al., "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin, Proc. Natl. Acad. Sci. U.S. A 7:8781-8781-8784 (1990). Sodoson et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At its Glycosylation Site," BioTechnology 8:343-346 (1990). Goodwin et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At its Glycosylation Site," BioTechnology 8:343-346 (1990). Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein." Proc. Natl. Acad. Sci. U. S. A. 87:7380-84 (1990). Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus." J. Clin. Endocrinol. Matab. 58:535-43 (1984). Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vitro and In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Biol. Chem. 260:12214-18 (1985). Haluser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40: 1735-39 (1990). Haluser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurolo | | | | | Group Art Unit | | | | | | | |
| OTHER DOCUMENTS (Including Marthor, Title, Date, Pertinent Pages, Etc). OTHER DOCUMENTS (Including Marthor, Title, Date, Pertinent Pages, Etc). Evans, "The steroid and thyroid hormone receptor superfamily," Science 240:889-95 (1988). Frohman et al., "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer," Proc. Natl. Acad. Sci. U. S. A. 85:8988-9002 (1988). Satanaga et al., "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin, Proc. Natl. Acad. Sci. U.S. A. 87:8781-8784 (1990). Soddwin et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At its Glycosylation Site," BioTechnology 8:343-346 (1990). Goodwin et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At its Glycosylation Site," BioTechnology 8:343-346 (1990). Goodwin et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At its Glycosylation Site," BioTechnology 8:343-346 (1990). Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein." Proc. Natl. Acad. Sci. U. S. A. 87:7380-84 (1990). Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus." J. Clin. Endocrinol. Metab. 58:535-43 (1984). Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Bio. Chem. 260:12214-18 (1985). Hatsey et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). Hatsey et al., "Cytokine accumulations in CSF of multiple | - 1 | (use | | essary 27 | Examiner Name | | | | | | | |
| Evans, "The steroid and thyroid hormone receptor superfamily," Science 240:889-95 (1988). Frohman et al., "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer," Proc. Natl. Acad. Sci. U. S. A. 85:8988-9002 (1988). Satisfied and the service of t | Sheet | ~~~ | | or \@ 9 | Attorney Docket No. | 98,385-1 | | | | | | |
| Frohman et al., "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer," <i>Proc. Natl. Acad. Sci. U. S. A.</i> 85:8998-9002 (1988). 83 Gatanaga et al., "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin, <i>Proc. Natl. Acad. Sci. USA 87:8781-8784</i> (1990). 84 Goodson et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At Its Glycosylation Site," BioTechnology 8:343-346 (1990). 85 Goodwin et al., "Molecular cloning and expression of the type 1 and type 2 murine receptors for tumor necrosis factor," <i>Mol. Cell. Biol.</i> 11:3020-26 (1991). 86 Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein," <i>Proc. Natl. Acad. Sci. U. S. A.</i> 87:7380-84 (1990). 87 Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," <i>J. Clin. Endocrinol. Metab.</i> 58:535-43 (1984). 88 Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) 89 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," <i>J. Biol. Chem.</i> 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," <i>Science</i> 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," <i>Neurology</i> 40:1735-39 (1990). 92 Hellecatal., "Complimentany, ONA, Cologing, of acceptor-for-Mamorenoerosis-factor-gened-demensis-factor-when form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93 Hi | 01 | SAL | OTHER DOCUM | ENTS (Including Author | or, Title, Date, Pertinent Pages, | Etc). | | | | | | |
| gene-specific oligonucleotide primer," Proc. Natl. Acad. Sci. U. S. A. 85:8998-9002 (1988). Gatanaga et al., "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin, Proc. Natl. Acad. Sci. USA 87:8781-8784 (1990). Goodson et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At Its Glycosylation Site," BioTechnology 8:343-346 (1990). Goodwin et al., "Molecular cloning and expression of the type 1 and type 2 murine receptors for tumor necrosis factor," Mol. Cell. Biol. 11:3020-26 (1991). Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein," Proc. Natl. Acad. Sci. U. S. A. 87:7380-84 (1990). Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," J. Clin. Endocrinol. Metab. 58:535-43 (1984). Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Biol. Chem. 260:12214-18 (1985). Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," Science 244:551-56 (1989). Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). Hauser et al., "Nolecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). | | 3 | Evans, "The steroid | and thyroid hormone | receptor superfamily," Science | e 240:889-95 (1988). | | | | | | |
| Gatanaga et al., "Purification and characterization of an inhibitor (soluble tumor necrosis factor receptor) for tumor necrosis factor and lymphotoxin, Proc. Natl. Acad. Sci. USA 87:8781-8784 (1990). 84 Goodson et al., "Site-Directed Pegylation of Recombinant Interleukin-2 At Its Glycosylation Site," BioTechnology 8:343-346 (1990). 85 Goodwin et al., "Molecular cloning and expression of the type 1 and type 2 murine receptors for tumor necrosis factor," Mol. Cell. Biol. 11:3020-26 (1991). 86 Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein," Proc. Natl. Acad. Sci. U. S. A. 87:7380-84 (1990). 87 Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," J. Clin. Endocrinol. Metab. 58:535-43 (1984). 88 Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coll," J. Cell Biochem. Suppl. 15F:113(1991) 89 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Biol. Chem. 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," Science 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). 92 Heller et al., "Complimentary DNA closing of a teoepter-fer-tumereneorosis-fector-and-demogration-science-shed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93 Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990 | ADEN. | AFE CON SEC | | | | | | | | | | |
| BioTechnology 8:343-346 (1990). 85 Goodwin et al., "Molecular cloning and expression of the type 1 and type 2 murine receptors for tumor necrosis factor," <i>Mol. Cell. Biol.</i> 11:3020-26 (1991). 86 Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein," <i>Proc. Natl. Acad. Sci. U. S. A.</i> 87:7380-84 (1990). 87 Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," <i>J. Clin. Endocrinol. Metab.</i> 58:535-43 (1984). 88 Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) 89 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," <i>J. Biol. Chem.</i> 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," <i>Science</i> 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," <i>Neurology</i> 40:1735-39 (1990). 92 Heller et al., "Complimentary DNA cloping of a tecepter for Human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," <i>DNA Cell Biol.</i> 9:705-15 (1990). 93 Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," <i>J. Exp. Med.</i> 170:607-12 (1989). | | | | | | | | | | | | |
| necrosis factor," Mol. Cell. Biol. 11:3020-26 (1991). 6 Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein," Proc. Natl. Acad. Sci. U. S. A. 87:7380-84 (1990). 6 Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," J. Clin. Endocrinol. Metab. 58:535-43 (1984). 8 Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) 8 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Biol. Chem. 260:12214-18 (1985). 9 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," Science 244:551-56 (1989). 9 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). Hellecet al., "Complimentary DNA cloning of a teceptor for tumor necrosis factor shed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Sci. 1989. | | 84 | Goodson et al., "Site BioTechnology 8:34 | e-Directed Pegylation (3-346 (1990). | of Recombinant Interleukin-2 | At Its Glycosylation Site," | | | | | | |
| recombinant soluble TNF-binding protein," <i>Proc. Natl. Acad. Sci. U. S. A.</i> 87:7380-84 (1990). 87 Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," <i>J. Clin. Endocrinol. Metab.</i> 58:535-43 (1984). 88 Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) 89 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," <i>J. Biol. Chem.</i> 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," <i>Science</i> 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sciencis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," <i>Neurology</i> 40:1735-39 (1990). 92 Heller et al., "Cytokine accumulations in CSF of multiple sciences paddemens tratice observable of the receptor," <i>Proc. Natl. Acad. Sci. USA Vol. 87</i> , pp. 6151-6155, August 1990. 93 Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," <i>DNA Cell Biol.</i> 9:705-15 (1990). 94 Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," <i>J. Exp. Med.</i> 170:607-12 (1989). | | 85 | Goodwin et al., "Mole necrosis factor," Mol | Goodwin et al., "Molecular cloning and expression of the type 1 and type 2 murine receptors for tumor necrosis factor," <i>Mol. Cell. Biol.</i> 11:3020-26 (1991). | | | | | | | | |
| Grizzard et al., "Affinity-labeled somatomedin-C receptors and binding proteins from the human fetus," <i>J. Clin. Endocrinol. Metab.</i> 58:535-43 (1984). Hale et al., "Cytokines and Their Receptors: From Clonal to Clinical Investigation, Demonstration of In Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," <i>J. Biol. Chem.</i> 260:12214-18 (1985). Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," <i>Science</i> 244:551-56 (1989). Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," <i>Neurology</i> 40:1735-39 (1990). Heller et al., "Complimentary DNA cloning of a teceptor-for-tumor-necrosis factor-and-demonstration-of-asshed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," <i>DNA Cell Biol.</i> 9:705-15 (1990). Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," <i>J. Exp. Med.</i> 170:607-12 (1989). | | 86 | Gray et al., "Cloning recombinant soluble | of human tumor necre TNF-binding protein, | osis factor (TNF) receptor cDI | NA and expression of A. 87:7380-84 (1990). | | | | | | |
| Vitro and In Vivo Efficacy of Two Biologically Active Human Soluble TNF Receptors Expressed in E. Coli," J. Cell Biochem. Suppl. 15F:113(1991) 89 Hass et al., "Characterization of specific high affinity receptors for human tumor necrosis factor on mouse fibroblasts," J. Biol. Chem. 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," Science 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). 92 Heller et al., "Complimentary DNA cloping of a receptor for tumor necrosis factor and demonstration of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93 Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). 94 Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). 95 Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, | | 87 | Grizzard et al., "Affin | ity-labeled somatome | din-C receptors and binding p | | | | | | | |
| fibroblasts," J. Biol. Chem. 260:12214-18 (1985). 90 Hatakeyama et al., "Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's," Science 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). 92 Heller et al., "Complimentary DNA cloning of a receptor-for-tumor-necrosis-factor-and-demonstration-of-asshed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93 Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis-factor receptor chain (p60) and its soluble derivative, tumor necrosis-factor-binding protein," DNA Cell Biol. 9:705-15 (1990). 94 Hofman et al., "Tumor necrosis-factor identified in multiple sclerosis-brain," J. Exp. Med. 170:607-12 (1989). 95 Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scit. 1989 | | 88 | Vitro and In Vivo Effi | cacy of Two Biologica | illy Active Human Soluble TNI | estigation, Demonstration of In Receptors Expressed in E. | | | | | | |
| human alpha and beta chain cDNA's," Science 244:551-56 (1989). 91 Hauser et al., "Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). 92 Heller et al., "Complimentary DNA closing of a receptor fortumor necrosis factor and demonstration of a shed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93 Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). 94 Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). 95 Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scat., 1989 | | 89 | Hass et al., "Charact fibroblasts," J. Biol. (| terization of specific h Chem. 260:12214-18 | igh affinity receptors for huma (1985). | n turnor necrosis factor on mouse | | | | | | |
| interleukin-1 and tumor necrosis factor but not interleukin-6," Neurology 40:1735-39 (1990). 92. Heller et al., "Complimentary DNA cloning of a receptor for tumor necrosis factor and demonstration of shed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. 93. Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). 94. Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). 95. Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scat., 1989 | | 90 | Hatakeyama et al., "I human alpha and be | Interleukin-2 receptor ta chain cDNA's," Sci | beta chain gene: generation of the description of t | of three receptor forms by cloned | | | | | | |
| shed form of the receptor," Proc. Natl. Acad. Sci. USA Vol. 87, pp. 6151-6155, August 1990. Himmler et al., "Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," J. Exp. Med. 170:607-12 (1989). Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scat. 1989 | V | 91 | Hauser et al., "Cytok interleukin-1 and turn | ine accumulations in nor necrosis factor but | CSF of multiple sclerosis patie t not interleukin-6," Neurology | ents: frequent detection of 40:1735-39 (1990). | | | | | | |
| (p60) and its soluble derivative, tumor necrosis factor-binding protein," <i>DNA Cell Biol.</i> 9:705-15 (1990). Hofman et al., "Tumor necrosis factor identified in multiple sclerosis brain," <i>J. Exp. Med.</i> 170:607-12 (1989). Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scat., 1989 | | 92 | Heller et al. "Compli shed form of the rec | mentary <u>DNA cloning</u> eptor," Proc. Natl. Aca | of_a.ceepter₅fer4trnereneores id. Sci. USA Vol. 87, pp. 6151 | sissfactorandsdemonstation, of a war- | | | | | | |
| (1989). 95 Hohmann, H.P. et al., J. Biol. Chem. 264:14927-14934, Scat. 1989 | EA | 93 | Himmler et al., "Mole (p60) and its soluble | cular cloning and exp derivative, tumor nec | ssion of human and rat tumor necrosis factor receptor chain sis factor-binding protein," DNA Cell Biol. 9:705-15 (1990). | | | | | | | |
| | | 94 | Hofman et al., "Tumo (1989). | or necrosis factor iden | tified in multiple sclerosis brai | n," J. Exp. Med. 170:607-12 | | | | | | |
| | | 95 | Hohmann, H.P. et al. | ., J. Biol. Chem. 264:1 | 4927-14934, Seet. | 1989 | | | | | | |
| Immunol. Lett. 12:217-224(1986). | 1 | 96 | Israel et al. "Binding | of Human TNF-alpha | | | | | | | | |

Burden Hour Statement: This form is estimated to take 2.0 hours to complete. Time will vary depending upon the needs of the Individual case. Any comments on the amount of time you are required to compete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC. 20231

^{*}EXAMINER: Initial If reference considered, whether or not citation is in conformance with MPEP 809. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

¹ Unique citation designation number. 2 Applicant is to place a check mark here if English translation is attached.

| ibstitute for | form 1449A/ | рто 📠 | Complete if Known | | | | | |
|---|---|---|---|--|--|--|--|--|
| | | - | Application No. | 09/898,234 | | | | |
| | INFOR | MATION DISCLOSURE | Filing Date: | 7-3-01 | | | | |
| | | MENT BY APPLICANTO PE | First Named Inventor | Hauptmann et al., | | | | |
| | | _ | Group Art Unit | | | | | |
| / | O(PRE | 5 . * HA 1 1 MINISTER | Examiner Name | | | | | |
| eet [| DIN 23 VI 3 | | Attomey Docket No. | 98,385-1 | | | | |
| MA. | | OTHER DOCUMENTS (Including Audior, | Title, Date, Pertinent Pages, | , Etc). | | | | |
| A | Manne | Seckinger et al., "A Human Inhibitor of Tumo | or Necrosis Factor Alpha," J | l. Exp. Med. 167:1511-1516. | | | | |
| | 128 | Seckinger et al., "A Urine Inhibitor of Interleu Tumor Necrosis Factor a," J. Immunol. 139(| | nterleuking 1 and B But Not | | | | |
| | 129 | Seckinger et al., "A Urine Inhibitor of Interleu Tumor Necrosis Factor a," J. Immunol. 139(| | nterleuking 1 and B But Not | | | | |
| | 130 | Selmaj et al., "Prevention of Chr-eae with Soluble TNF Receptor P55," J. Neuramman 1033 | | | | | | |
| | 131 | Selmaj and Raine, "Anti-Tumor Necrosis Factor Therapy Abrogates Autoimmune Demyelination, Annals of Neurology Vol 30 No 5 November 1991. | | | | | | |
| | 132 | Shimuzu et al., PNAS USA, 80:2112-2116 (| 1983). | | | | | |
| Smith et al., "A Receptor for Tumor Necrosis Factor Defines an Unusual Family Cellu Proteins, Reports, 1019-1023. Science, May 1990. | | | | | | | | |
| - | 134- | -Smithet-ala-Science-248:1019-1023(1990) | NAME OF THE OWNER, OWNER, OWNER, OWNER, OWNER, OWNER, | THE RESERVE THE PROPERTY OF TH | | | | |
| FA | actor block its binding to cell- | | | | | | | |
| | 136 | surface receptor," <i>Proc. Natl. Acad. Sci. U. S. A.</i> 84:8829-33 (1987). Spinas et al., "Induction of plasma inhibitors of interleukin 1 and TNF-alpha activity by endotoxin administration to normal humans," <i>Am. J. Physiol.</i> 259(5 Pt 2):R993-7 (1990). | | | | | | |
| | 137 | Stauber and Aggarwal, "Characterization an lymphotoxin (tumor necrosis factor-beta) on Chem. 264:3573-76 (1989). | d affinity cross-linking of red | ceptors for human recombinant | | | | |
| | 138 | Stauber. G. et al., J. Biol. Chem. 263(35):19 | 098-19104. Dec. | 19 88 | | | | |
| | 139 | Suffys et al., "Involvement of a serine protea Biochem. 178:257-65 (1988). | | <u> </u> | | | | |
| | 140 | Suggs et al., "Use of synthetic oligonucleotic sequences for human beta 2-microglobulin," | des as hybridization probes: Proc. Natl. Acad. Sci. U. S | isolation of cloned cDNA . A. 78:6613-17 (1981). | | | | |
| | 141 | The Cytokine Factsbook pp. 244-46 (Callard | | | | | | |
| | 142 | Tracey et al., "Cachectin/tumor necrosis fact Med. 167:1211-27 (1988). | tor induces cachexia, anem | ia, and inflammation," <i>J. Exp.</i> | | | | |
| | Tracey et al., "Metabolic effects of cachectin/tumor necrosis factor are modified by site of product Cachectin/tumor necrosis factor-secreting tumor in skeletal muscle induces chronic cachexia, wimplantation in brain induces predominantly acute anorexia," J. Clin. Invest. 86:2014-24 (1990). | | | | | | | |

Clear B.0 Hara 1/21/03

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

¹ Unique citation designation number. ² Applicant is to place a check mark here if English translation is attached.

| | Substitute for form 144 | 9A/PTO | | Complete if Known | | |
|------|-------------------------|------------------|-----------------|-------------------|----------------------|------------|
| < | PE | | | | Application No. | 09/898,234 |
| J | INFOR | RMATION | DISCLOS | URE | Filing Date: | 7-3-01 |
| AUE | 0 1 2003S副ATI | EMENT BY | / APPLIC | ANT | First Named Inventor | Hauptmann |
| ٥ | .0/ | | . , | ,,,,,,, | Group Art Unit | 1646 |
| VEN. | & TRADEMARK (U | se as many sheet | ts as necessary |) | Examiner Name | E. O'Hara |
| | Sheet | 1 | of | 2 | Attorney Docket No. | 98-385-I |

| | | | | U.S. PATENT DOCUMENT | 'S | |
|--------------------|--------------|----------------------|--------------------------------------|---|------------------------------|---|
| Examiner Initials* | Cite No.1 | U.S. Patent Document | | | Date of Publication of | Pages, Columns, Lines |
| | | Number | Kind Code ² (if known) | Name of Patentee or Applicant of Cited Document | Cited Document MM-DD-YYYY | Where Relevant Passages or Figures Appear |
| ENA | | 6,294,352 | B1 | Hauptmann et al. | 09-25-2001 | |
| | | | | | | |

| | FOREIGN PATENT DOCUMENTS | | | | | | | | | |
|-----------|--------------------------|------------------------------|---------------------|--|--|------------------------------|---|----|--|--|
| Examiner | Cite | Cite Foreign Patent Document | | | | Date of Publication of | Pages, Columns, Lines | | | |
| Initials* | No.¹ | Office ³ | Number ⁴ | Number ⁴ Kind Code ⁵ Applicant of Cited Document | | Cited Document MM-DD-YYYY | Where Relevant Passages or Figures Appear | T⁵ | | |
| EM | J | DE | 3913101.7 | A1 | Boehringer Ingelheim International GmbH | 04/29/1989 | | | | |
| | | | | | | | | | | |

| | | OTHER DOCUMENTS NON PATENT LITERATURE DOCUMENTS | |
|-----------------------|--------------------------|---|----------------|
| Examiner Initials* | Cite No. ¹ | Include name of author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published | T ² |
| Ext | 7 | HELLER et al., "Complementary DNA cloning of a receptor for tumor necrosis factor and demonstration of a shed form of the receptor", Proc. Natl. Acad. Sci. 87: 6151-55 (1990). | |
| | 4 | PEPPEL et al., "A Tumor Necrosis Factor (TNF) Receptor-lgG Heavy Chain Chimeric Protein as a Bivalent Antagonist of TNF Activity," J. Exp. Med. 174:1483-89 (1991). | |
| | J | POWELL et al., "Lymphotoxin and tumor necrosis factor-alpha production by myelin basic protein specific T-cell clones correlates with encephalitogenicity," Int. Immunol. 2:539-44 (1990). | |
| | 7 | POWELL et al., "The Role of Lymphotoxin and TNF in Demyelinating Diseases of the CNS," Tumor Necrosis Factors: The Molecules and Their Emerging Role in Medicine. pgs 355-69. 199. | 2 |
| | | RHEIN et al., "Another Sepsis Drug Down-Immunex TNF Receptor," Biotechnology Newswatch, pg. 1,3 (Monday, October 4, 1991) | |
| | 7 | VILCEK et al., "Tumor Necrosis Factor: Receptor Binding and Mitogenic Action in Fibroblasts," J. Cell. Physiol. 5:57-61 (1987). | |
| A | 1 | SELMAJ et al., "Tumor Necrosis Factor Mediates Myelin and Oligodendrocyte Damage In Vitro," Ann. Neurol. 23(4):339-46 (1988). | |

| Examiner | En RAII | Date / / / |
|-----------|----------------|---------------------|
| Signature | Ciller Di Hara | Considered 10/13/83 |

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 809. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Burden Hour Statement: This form is estimated to take 2.0 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to compete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC. 20231

The Gran

¹ Unique citation designation number. ² See attached Kinds of U.S. Patent Documents. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST.3). ⁴ For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial number of the patent document. ⁵ Kind of document by the appropriate symbols as indicated on the document under WIPO Standard ST.16 if possible. ⁶ Applicant is to place a check mark here if English translation is attached.